

Low serum 25-hydroxyvitamin D is associated with higher risk of frequent headache in middle-aged and older men

Virtanen J., Giniatullin R., Mäntyselkä P., Voutilainen S., Nurmi T., Mursu J., Kauhanen J., Tuomainen T.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© The Author(s) 2017. Vitamin D has been suggested to have a role in various neurovascular diseases, but the data regarding headache is inconclusive. Our aim was to investigate the associations between serum 25-hydroxyvitamin D [25(OH)D], a marker for vitamin D status, and risk of frequent headache. The study population consisted of 2601 men from the population-based Kuopio Ischaemic Heart Disease Risk Factor Study (KIHD) from eastern Finland, aged 42–60 years in 1984–1989. The cross-sectional associations with prevalence of self-reported frequent headache (defined as weekly or daily headaches) were estimated with multivariable-adjusted odds ratios. The average serum 25(OH) concentration was 43.4 nmol/L (SD 18.9, min–max 7.8–136.1 nmol/L). A total of 250 men (9.6%) reported frequent headache. The average serum 25(OH)D concentration among those with frequent headache was 38.3 nmol/L (SD 18.8) and 43.9 nmol/L (SD 18.9) among those without frequent headache, after adjustment for age and year and month of blood draw (P for difference < 0.001). After multivariable adjustments, those in the lowest vs. the highest serum 25(OH)D quartile had 113% (95% CI 42, 218%; P for trend < 0.001) higher odds for frequent headache. In conclusion, low serum 25(OH)D concentration was associated with markedly higher risk of frequent headache in men.

<http://dx.doi.org/10.1038/srep39697>

References

- [1] Smitherman, T. A., Burch, R., Sheikh, H. & Loder, E. The prevalence, impact, and treatment of migraine and severe headaches in the United States: a review of statistics from national surveillance studies. *Headache* 53, 427–436 (2013).
- [2] Westergaard, M. L., Hansen, E. H., Glumer, C., Olesen, J. & Jensen, R. H. Definitions of medication-overuse headache in population-based studies and their implications on prevalence estimates: a systematic review. *Cephalalgia* 34, 409–425 (2014).
- [3] Chiang, C. C., Schwedt, T. J., Wang, S. J. & Dodick, D. W. Treatment of medication-overuse headache: A systematic review. *Cephalalgia* 36, 371–386 (2016).
- [4] Munksgaard, S. B. & Jensen, R. H. Medication overuse headache. *Headache* 54, 1251–1257 (2014).
- [5] Buell, J. S. & Dawson-Hughes, B. Vitamin D and neurocognitive dysfunction: preventing "D"ecline? *Mol. Aspects Med.* 29, 415–422 (2008).
- [6] Prabhakar, P., Majumdar, V., Kulkarni, G. B. & Christopher, R. Genetic variants of vitamin D receptor and susceptibility to ischemic stroke. *Biochem. Biophys. Res. Commun.* 456, 631–636 (2015).
- [7] Prakash, S. et al. The prevalence of headache may be related with the latitude: a possible role of Vitamin D insufficiency? *J. Headache Pain* 11, 301–307 (2010).

- [8] Motaghi, M. et al. Relationship between vitamin D receptor gene polymorphisms and migraine without aura in an Iranian population. *Biomed. Res. Int.* 2013, 351942 (2013).
- [9] Kjaergaard, M., Eggen, A. E., Mathiesen, E. B. & Jorde, R. Association between headache and serum 25-hydroxyvitamin D: the Tromsø Study. *Headache* 52, 1499-1505 (2012).
- [10] Zandifar, A. et al. Vitamin D status in migraine patients: a case-control study. *Biomed. Res. Int.* 2014, 514782 (2014).
- [11] Knutsen, K. V. , Brekke, M., Gjelstad, S. & Lagerlov, P. Vitamin D status in patients with musculoskeletal pain, fatigue and headache: a cross-sectional descriptive study in a multi-ethnic general practice in Norway. *Scand. J. Prim. Health Care* 28, 166-171 (2010).
- [12] Tys-Jacobs, S. Vitamin D and calcium in menstrual migraine. *Headache* 34, 544-546 (1994).
- [13] Tys-Jacobs, S. Alleviation of migraines with therapeutic vitamin D and calcium. *Headache* 34, 590-592 (1994).
- [14] Prakash, S., Makwana, P. & Rathore, C. Vitamin D deficiency mimicking chronic tension-type headache in children. *BMJ Case Rep.* 2016, 10.1136/bcr-2015-213833 (2016).
- [15] Knutsen, K. V. et al. Effect of vitamin D on musculoskeletal pain and headache: A randomized, double-blind, placebo-controlled trial among adult ethnic minorities in Norway. *Pain* 155, 2591-2598 (2014).
- [16] Mottaghi, T., Askari, G., Khorvash, F. & Maracy, M. R. Effect of Vitamin D supplementation on symptoms and C-reactive protein in migraine patients. *J. Res. Med. Sci.* 20, 477-482 (2015).
- [17] Buettner, C. et al. Simvastatin and vitamin D for migraine prevention: a randomized controlled trial. *Ann. Neurol.* 78, 970-981 (2015).
- [18] Buettner, C. & Burstein, R. Association of statin use and risk for severe headache or migraine by serum vitamin D status: a cross-sectional population-based study. *Cephalalgia* 35, 757-766 (2015).
- [19] Salonen, J. T. Is there a continuing need for longitudinal epidemiologic research? The Kuopio Ischaemic Heart Disease Risk Factor Study. *Ann. Clin. Res.* 20, 46-50 (1988).
- [20] Salonen, J. T. et al. High stored iron levels are associated with excess risk of myocardial infarction in eastern Finnish men. *Circulation* 86, 803-811 (1992).
- [21] Lakka, T. A. et al. Relation of leisure-time physical activity and cardiorespiratory fitness to the risk of acute myocardial infarction. *N. Engl. J. Med.* 330, 1549-1554 (1994).
- [22] Nurmi, T., Tuomainen, T. P. , Virtanen, J., Mursu, J. & Voutilainen, S. High-performance liquid chromatography and coulometric electrode array detector in serum 25-hydroxyvitamin D(3) and 25-hydroxyvitamin D(2) analyses. *Anal. Biochem.* 435, 1-9 (2013).
- [23] Shipton, E. E. & Shipton, E. A. Vitamin D deficiency and pain: Clinical evidence of low levels of vitamin D and supplementation in chronic pain states. *Pain Ther.* 4, 67-87 (2015).
- [24] Carlberg, C. Genome-wide (over)view on the actions of vitamin D. *Front. Physiol.* 5, 167 (2014).
- [25] Yu, C., Fedoric, B., Anderson, P. H., Lopez, A. F. & Grimbaldston, M. A. Vitamin D(3) signalling to mast cells: A new regulatory axis. *Int. J. Biochem. Cell Biol.* 43, 41-46 (2011).
- [26] Tague, S. E. & Smith, P. G. Vitamin D receptor and enzyme expression in dorsal root ganglia of adult female rats: modulation by ovarian hormones. *J. Chem. Neuroanat.* 41, 1-12 (2011).
- [27] Waltman, N. L., Ott, C. D., Twiss, J. J., Gross, G. J. & Lindsey, A. M. Vitamin D insufficiency and musculoskeletal symptoms in breast cancer survivors on aromatase inhibitor therapy. *Cancer Nurs.* 32, 143-150 (2009).
- [28] Prakash, S. & Shah, N. D. Chronic tension-type headache with vitamin D deficiency: casual or causal association? *Headache* 49, 1214-1222 (2009).
- [29] Prakash, S., Kumar, M., Belani, P. , Susvirkar, A. & Ahuja, S. Interrelationships between chronic tension-type headache, musculoskeletal pain, and vitamin D deficiency: Is osteomalacia responsible for both headache and musculoskeletal pain? *Ann. Indian. Acad. Neurol.* 16, 650-658 (2013).
- [30] Giniatullin, R., Bart, G. & Tavi, P. Complex role of peroxisome proliferator activator receptors (PPARs) in nociception. *Scand. J. Pain* 9, 70-71 (2015).
- [31] Lajdova, I., Chorvat, D., Jr & Chorvatova, A. Rapid effects of 1 α ,25(OH) $_2$ D $_3$ in resting human peripheral blood mononuclear cells. *Eur. J. Pharmacol.* 586, 14-23 (2008).
- [32] Inoue, K. & Tsuda, M. Purinergic systems, neuropathic pain and the role of microglia. *Exp. Neurol.* 234, 293-301 (2012).